Home Journals Books Collections Resources Services Subscribe

informa



Which content access option is best for you?

Contact us to learn more

Clinical Toxicology

Login Register Athens Shibboleth

Advanced Search

All Books and Journals

All Issues

Current Issue

Early Online

Issue TOC | Previous Article | Next Article SHARE

Aims & Scope

Editorial Board

Instructions for Authors

PDF (491 KB)

Reprints

Permissions

PDF Plus (554 KB)



personal

subscription to

www.clintox.org/

Critical Care

Acute kidney injury associated with smoking synthetic cannabinoid

August 2014, Vol. 52, No. 7, Pages 664-673 (doi:10.3109/15563650.2014.932365)

G. L. Buser, ^{1,2}R. R. Gerona, ³B. Z. Horowitz, ⁴K. P. Vian, ⁵M. L. Troxell, ⁶R. G. Hendrickson, ⁴D. C. Houghton, ⁶D. Rozansky, ⁷S. W. Su, ⁸ and R. F. Leman¹

¹Acute and Communicable Disease Prevention, Oregon Public Health Division, Oregon Health Authority, Portland, OR, USA

²Epidemic Intelligence Service, Centers for Disease Control and Prevention, Atlanta, GA, USA

³Department of Laboratory Medicine, University of California, San Francisco, CA, USA

⁴Oregon Poison Center, Oregon Health and Science University, Portland, OR, USA

⁵Communicable Disease and Immunization Programs, Douglas County Public Health, Roseburg, OR LISA

⁶Department of Pathology, Oregon Health and Science University, Portland, OR, USA

⁷Department of Nephrology, Oregon Health and Science University, Portland, OR, USA

⁸Department of Pediatric Nephrology, Legacy Randall Children's Hospital, Portland, OR, USA Address correspondence to Genevieve L. Buser MDCM, MSHP, Acute and Communicable Disease Prevention, Oregon Public Health Division, Oregon Health Authority, 800 NE Oregon Street, Suite 772, Portland, OR 97232, USA. Tel: +(971) 673-1111. Fax: +(971) 673-1100. E-mail: genevieve.buser@gmail.com



Request a call back



Publish with us



Subscribe today

Informa Healthcare Reading Room

Read the latest

Abstract

Context and objectives. Synthetic cannabinoids are illegal drugs of abuse known to cause adverse neurologic and sympathomimetic effects. They are an emerging health risk: 11% of high school seniors reported smoking them during the previous 12 months. We describe the epidemiology of a toxicologic syndrome of acute kidney injury associated with synthetic cannabinoids, review the toxicologic and public health investigation of the cluster, and describe clinical implications of the cluster investigation. Materials and methods. Case series of nine patients affected by the toxicologic syndrome in Oregon and southwestern Washington during May-October 2012. Cases were defined as acute kidney injury (creatinine > 1.3 mg/dL) among persons aged 13-40 years without known renal disease who reported smoking synthetic cannabinoids. Toxicology laboratories used liquid chromatography and time-of-flight mass spectrometry to test clinical and product specimens for synthetic cannabinoids, their metabolites, and known nephrotoxins. Public health alerts informed clinicians, law enforcement, and the community about the cluster and the need to be alert for toxidromes associated with emerging drugs of abuse. Results. Patients were males aged 15-27 years (median, 18 years), with intense nausea and flank or abdominal pain, and included two sets of siblings. Peak creatinine levels were 2.6-17.7 mg/dL (median, 6.6 mg/dL). All patients were hospitalized; one required dialysis; none died. No alternate causes of acute kidney injury or

Quick Links



Add to Favourites

Download Citations

Email TOC Alert

Related Articles RSS TOC Alerts Email Citation Alert

Purchase



Informa Healthcare offers a unique range of publishing formats.

Podcasts

- *** CTX Interview with Dr. Priyanka Vakkalanka
- CTX Interview with Dr. Suzanne Doyon
- CTX Interview with Dr. Jess Benson



Users Also Read

Comprehensive analysis of "bath salts" purchased from California stores and the internet

free and Open Access articles from our journals



Click here to find out more nephrotoxins were identified. Patients reported easily purchasing synthetic cannabinoids at convenience, tobacco, and adult bookstores. One clinical and 2 product samples contained evidence of a novel synthetic cannabinoid, XLR-11 ([1-(5-fluoropentyl)-1H-indol-3-yl](2,2,3,3-tetramethylcyclopropyl)methanone). *Discussion and conclusion*. Whether caused by direct toxicity, genetic predisposition, or an as-yet unidentified nephrotoxin, this association between synthetic cannabinoid exposure and acute kidney injury reinforces the need for vigilance to detect new toxicologic syndromes associated with emerging drugs of abuse. Liquid chromatography and time-of-flight mass spectrometry are useful tools in determining the active ingredients in these evolving products and evaluating them for toxic contaminants.

'Crazy Monkey' Poisons Man and Dog: Human and canine seizures due to PB-22, a novel synthetic cannabinoid

Effects of hydroxocobalamin on carboxyhemoglobin measured under physiologic and pathologic conditions

More »

Keywords

Epidemiology, Electronspray ionization, Mass, Public health, Substance-related disorders, Spectrometry

HTML | PDF (491 KB) | PDF Plus (554 KB) - What is PDF Plus? |

 Journals
 Books
 About Us
 Careers at Informa

 View All Journals
 View All Books
 Privacy policy
 Advertise

Copyright © 2014 Informa Plc. All rights reserved.

Online Books

This site is owned and operated by Informa plc ("Informa") whose registered office is Mortimer House, 37-41 Mortimer Street, London, W1T 3JH. Registered in England and Wales. Number 309067. UK VAT Group: GB 365 4626 3

Terms & Conditions

Subscribe