


LC/MS/MS Analysis of Vitamin D Metabolites on Ascentis® Express F5

The metabolic role and health implications of the various vitamin D isoforms are of current clinical interest. Therefore, it is important to have an analytical method that will resolve all of the known isoforms with necessary sensitivity and specificity. This application shows an Ascentis Express F5 provided rapid resolution of the isobaric vitamin D metabolites. MS detection provides the necessary secondary resolution. Fluka LC-MS Ultra CHROMASOLV solvents were used to supply low background interference and low particulate contaminants for robust, trouble-free operation. Cerilliant CRMs provided reliable identification and quantification.

market focus	Clinical; Vitamins, Nutraceuticals, and Natural Products
column	Ascentis Express F5, 15 cm x 2.1 mm I.D., 2.7 µm particles (53571-U)
mobile phase	[A] 5 mM ammonium formate water; [B] methanol; (25:75, A:B)
flow rate	0.4 mL/min
pressure	5178 psi (357 bar)
column temp.	40 °C
detector	ESI(+), 100-1000 m/z
injection	2 µL, each compound 300 ng/mL in 40:60 (water:methanol)
Application No.	G005878



Related Products

analytical column

Ascentis® Express F5, 2.7 Micron HPLC Column ([Supelco 53571-U](#))

mobile phase component

Ammonium formate ([Fluka 14266](#))

Methanol ([Fluka 14262](#))

Water ([Fluka 14263](#))

standard

1 α ,25-Dihydroxyvitamin D₂ solution ([Cerilliant H-090](#))

1 α ,25-Dihydroxyvitamin D₃ solution ([Cerilliant H-089](#))

1 α -Hydroxyvitamin D₃ ([Sigma 17946](#))

3-*epi*-25-Hydroxyvitamin D₃ ([Aldrich 705993](#))

25-Hydroxyvitamin D₂ solution ([Cerilliant H-073](#))

25-Hydroxyvitamin D₃ solution ([Cerilliant H-083](#))

3-*epi*-25-Hydroxyvitamin D₂ solution ([Aldrich 753556](#))